



Short Communication

Emotional reactions to alcohol-related words: Differences between low- and high-risk drinkers

Carlos Gantiva^{a,*}, Rafael Delgado^b, Tania Romo-González^c^a University of San Buenaventura, Bogotá, Colombia^b University of Granada, Spain^c Veracruzana University, Mexico

HIGHLIGHTS

- Alcohol-related words are more appetitive for high-risk drinkers.
- High-risk drinkers perceive alcohol-related words as highly arousing.
- For alcohol-related words, high AUDIT scores are related to less dominance.

ARTICLE INFO

Available online 9 June 2015

Keywords:

Alcohol
Words
Emotion
Valence
Arousal

ABSTRACT

Introduction: Research that has examined responses to alcohol-related words in drinkers has mostly linked such responses to memory, attentional, and perceptual bias. However, studies of emotional processing in alcoholics have not received much attention. The main goal of the present study was to identify the features and differences of emotional responses to alcohol-related words in low- and high-risk drinkers.

Method: A total of 149 low-risk drinkers and 125 high-risk drinkers evaluated five alcohol-related words and 15 words from the Affective Norms for English Words in the dimensions of valence, arousal, and dominance using the Self-Assessment Manikin.

Results: The results indicated that high-risk drinkers evaluated alcohol-related words as more appetitive and arousing.

Conclusion: These results, together with findings in the attention and memory research literature, suggest that alcohol-related words can serve as conditioned cues in alcohol consumption.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Previous research has demonstrated the ability of words to prompt emotional reactions in humans (Bradley & Lang, 1999). These studies indicate that words that have appetitive and aversive content generate high arousal, and words that have neutral valence prompt low arousal. This pattern has been described across different countries (Kristensen, Gomes, Justo, & Vieira, 2011; Redondo, Fraga, Padrón, & Comesaña, 2007; Soares, Comesaña, Pinheiro, Simões, & Frade, 2012).

Previous studies have also demonstrated that words facilitate the recognition of emotions in facial expressions (Gendron, Lindquist, Barsalou, & Barrett, 2012), modulate the activation of facial muscles that are sensitive to stimulus valence (Herbert, Deutsch, Sütterlin, Kübler, & Pauli, 2011), and allow the study of neurological processes that are associated with emotions (Herbert, Kissler, Junghöfer, Peyk, & Rockstroh, 2006). Words can also reliably capture an individual's

attention, especially those that depict sexual and aversive content (Aquino & Arnell, 2007; Arnell, Killman, & Fijavz, 2007), but the effect diminishes rapidly with aversive words (Harris & Pashler, 2004).

Research that has used words with alcoholic patients has focused on assessing attention, memory bias (Fridrici et al., 2014; Johnsen, Laberg, Cox, Vaksdal, & Hugdahl, 1994) and emotional reactions (Stormark, Laberg, Nordby, & Hugdahl, 2000). Nevertheless, studies on emotional responses to alcohol-related words have been inconclusive. For example, Stormark et al. (2000) reported that alcohol-dependent subjects exhibited a significantly greater skin conductance response and greater heart rate deceleration in response to alcohol-related words compared with neutral and aversive words; however, no differences were found between alcoholic and nonalcoholic participants. Additionally, this study did not use appetitive words, which may be pivotal when considering that social interaction and advertising associated with alcohol consumption are related to appetitive cues, which is reflected in verbal and written language (van Zyl & Meiselman, 2015). Furthermore, studies on emotional responses to alcohol-related words have only been performed with alcoholic patients. Differences between low- and

* Corresponding author at: Cr 8 H # 172-20, Bogotá, Colombia. Tel./fax: + 57 1 6671090. E-mail addresses: cgantiva@usbog.edu.co, cgantiva@gmail.com (C. Gantiva).

high-risk drinkers have not been examined. Such drinkers represent the typology of the largest proportion of drinkers in the population.

To address this issue, the present study sought to identify the features and differences of emotional responses to alcohol-related words in low- and high-risk drinkers and make comparisons with emotional responses to affective and neutral words.

2. Methods

2.1. Participants

A total of 274 university students (139 females) voluntarily participated in the study. The participants were distributed into two groups according to scores on the AUDIT (WHO, 2001): low-risk drinkers ($n = 149$; scores between 1 and 7; $M = 3.89$, $SD = 1.85$) and high-risk drinkers ($n = 125$; scores between 8 and 19; $M = 11.56$, $SD = 3.54$; $t(272) = -22.92$, $p < .0001$). No significant differences were found between groups in age, percent of men and women, and years of drinking (all $p > .49$). A 3-year history of alcohol consumption was used as the inclusion criterion. The study was approved by the University of San Buenaventura Review Board and all of the subjects signed written informed consent forms.

2.2. Stimuli

Fifteen words (five pleasant, five neutral, five unpleasant) were selected from the Affective Norms for English Words (ANEW) (Bradley & Lang, 1999), which were translated to Spanish by Redondo et al. (2007). Five alcohol-related words¹ were also included. For the selected ANEW words significant differences were found between word categories in the valence dimension (all $p < .0001$). For the arousal dimension, significant differences were found between affective words (pleasant and unpleasant) and neutral words (both $p = .001$).

2.3. Measures

The Self-Assessment Manikin (SAM; Bradley & Lang, 1994) is a non-verbal pictographic scale designed to assess a participant's feelings in three emotional dimensions: valence (pleasant vs. unpleasant), arousal (relaxed vs. activated), and dominance (feeling in control vs. feeling controlled). Each dimension is represented by humanoid figures that indicate different levels of intensity.

2.4. Procedure

In small groups of not more than 28 individuals, the participants were provided with booklets that contained the SAM scales to evaluate the words. The Spanish instructions were presented in a digital audio format. After instructions were given to the participants, three words were used as examples, one word from each of the affective categories (Friend, Cabinet, and Killer). The instructions and words were presented using a projector. The words were presented in white against a black background in the center of the screen in 80-pt Arial font. Each trial consisted of three parts: 6 s of word presentation, 15 s to rate the word using the SAM scale, and a 5 s intertrial interval. After the words were presented, the participants completed the AUDIT. Four different pseudo-randomized word presentation orders were prepared. Each order had the constraint of not presenting the same word category consecutively more than twice.

¹ Pleasant words: Kiss, Caress, Chocolate, Orgasm, and Treasure. Neutral words: Street, Basket, Square, Table, and Paper. Unpleasant words: Infection, Abuse, Dirty, Torture, and Rape. Alcohol-related words: Beer, Bar, Party, Drunk, and Drink.

2.5. Statistical analysis

Pearson's linear correlation was used to analyze correlations between valence and arousal, and valence and dominance. To examine the emotional processing of alcohol-related words and ANEW words in the two alcohol consumption groups, we ran separate mixed ANOVAs 2×4 for each emotional dimensions, with type of alcohol consumption as the between-subjects factor (low- and high-risk drinkers) and word category (pleasant, neutral, unpleasant, and alcohol-related) as the within-subjects factor. Post hoc analyses of the mean values were performed using paired multiple comparisons, adjusted with the Bonferroni correction. Finally, a Pearson correlation was used to analyze the association between AUDIT scores and SAM ratings for alcohol-related words. The level of significance was set at $p < .05$. All of the statistical analyses were performed using SPSS 20.0 software.

3. Results

3.1. Affective space

Fig. 1A and B illustrates the distribution of the 20 words in the two-dimensional affective space that was composed of the valence and arousal dimensions. In both groups, pleasant words were very appetitive and highly arousing. Neutral words had an intermediate valence and were slightly arousing. Unpleasant words were aversive but not highly arousing. Alcohol-related words had a different distribution for low- and high-risk drinkers. Low-risk drinkers evaluated them with an intermediate valence and arousal (Fig. 1A). By contrast, they were very appetitive and highly arousing for high-risk drinkers (Fig. 1B).

The quadratic correlation was positive and significant for low-risk drinkers ($R^2 = .96$, $p < .0001$) and high-risk drinkers ($R^2 = .94$, $p < .0001$), suggesting the influence of valence on arousal. For low-risk drinkers, Pearson's correlation between appetitive valence and arousal was positive and significant ($r = .97$, $p < .0001$, $R^2 = .94$) and negative and significant for aversive valence and arousal ($r = -.93$, $p < .0001$, $R^2 = .86$). Pearson's correlation was also positive and significant between appetitive valence and arousal for high-risk drinkers ($r = .94$, $p < .0001$, $R^2 = .88$), but it was negative and not significant between aversive valence and arousal ($r = -.14$, $p = .66$, $R^2 = .01$).

Fig. 1C and D shows the distribution of the 20 words in the two-dimensional affective space that was composed of the dominance and valence dimensions. The words were generally distributed similarly in both groups, but alcohol-related words were placed in a similar region as neutral words for low-risk drinkers (Fig. 1C). In contrast, in high-risk drinkers, words that depicted alcohol content were placed in a region that was closer to pleasant words (Fig. 1D). A positive and significant linear relationship was found in both groups (low-risk drinkers: $r = .88$, $p < .0001$, $R^2 = .77$; high-risk drinkers: $r = .86$, $p < .0001$, $R^2 = .73$).

3.2. Emotional dimensions

3.2.1. Valence

The ANOVA revealed a significant main effect of word category ($F_{3,516} = 732.59$, $p < .0001$, $\eta^2 = .79$). Pleasant words were the most appetitive and unpleasant words were the most aversive (all $p < .0001$). Alcohol-related words were perceived as more appetitive than unpleasant and neutral words but less appetitive than pleasant words (all $p < .0001$). A significant quadratic trend was obtained ($F_{1,172} = 1140.26$, $p < .0001$, $\eta^2 = .86$). A significant Word \times Alcohol consumption interaction was found ($F_{3,516} = 14.96$, $p < .0001$, $\eta^2 = .01$). High-risk drinkers perceived alcohol-related words as more appetitive than low-risk drinkers ($p < .001$), and low-risk drinkers evaluated neutral words with more valence than high-risk drinkers ($p < .001$) (Fig. 2A).




Fig. 1. Distribution of the 20 words in the two-dimensional affective space. (A) Valence and arousal in low-risk drinkers. (B) Valence and arousal in high-risk drinkers. (C) Valence and dominance in low-risk drinkers. (D) Valence and dominance in high-risk drinkers.

3.2.2. Arousal

The ANOVA revealed a significant main effect of word category ($F_{3,516} = 205.98, p < .0001, \eta^2 = .52$), indicating that the participants felt more arousal when reading the pleasant words (all $p < .0001$). Alcohol-related words were perceived as more arousing than unpleasant and neutral words and unpleasant words were evaluated as more arousing than neutral words (all $p < .03$). A significant quadratic trend was found ($F_{1,172} = 479.98, p < .0001, \eta^2 = .71$). A significant Word \times Alcohol consumption interaction was found ($F_{3,516} = 13.80, p < .0001, \eta^2 = .03$). Low-risk drinkers felt more arousal when reading

neutral and unpleasant words (both $p < .03$), and high-risk drinkers felt more arousal when reading alcohol-related words compared with low-risk drinkers ($p = .001$) (Fig. 2B).

3.2.3. Dominance

The ANOVA revealed a significant main effect of word category ($F_{3,516} = 103.49, p < .0001, \eta^2 = .37$). The participants felt more dominant (i.e., in control) when reading pleasant and alcohol-related words compared with neutral words (both $p < .001$). Furthermore, the participants felt less dominant when reading unpleasant words compared




Fig. 2. Emotional reactions to affective and alcohol-related words in low- and high-risk drinkers. (A) Valence dimension. (B) Arousal dimension. Bars are standard error of the mean. * $p < .05$.

with the other word categories (all $p < .0001$). A significant quadratic trend was obtained ($F_{1,172} = 133.20, p < .0001, \eta^2 = .43$).

3.3. Correlation analysis

For alcohol-related words, the subjects' AUDIT scores were positively and significantly correlated with SAM ratings in the affective dimensions of valence ($r = .55, p < .0001, R^2 = .30$) and arousal ($r = .50, p < .0001, R^2 = .25$). A significant negative correlation was found between AUDIT scores and the dominance scale ($r = -.16, p = .007, R^2 = .02$).

4. Discussion

The results showed that alcohol-related words were perceived as appetitive and highly arousing in high-risk drinkers. Low-risk drinkers evaluated them as moderately appetitive and slightly arousing.

Previous studies evaluated emotional responses to words that depicted different affective contents and found that words that were perceived with either appetitive or negative valence also generated high arousal, and words that were evaluated with intermediate valence generated low arousal (Bradley & Lang, 1999). However, the words that were perceived as aversive in the present study did not generate a high level of arousal. These results may be explained by the characteristics of the sample (i.e., unselected undergraduate participants). Evidence suggests that fear/threat words are rather weak emotional cues in the normal population compared with patients diagnosed with anxiety disorders (Williams, Mathews, & MacLeod, 1996). Previous research also showed that fear/threat words less powerfully prompt emotional responses compared with sexual words in the normal population (Arnell et al., 2007).

Concerning alcohol-related words, the present results showed that high-risk drinkers rated them as more appetitive and arousing compared with low-risk drinkers. These results may suggest that these words serve as alcohol-related cues that activate appetitive motivational systems in high-risk drinkers, which would suggest a greater likelihood of approach behaviors (Lang, 2010). These findings, together with attentional (Stormark et al., 2000), interpretation (Woud et al., 2014), and memory (Fridrici et al., 2014) bias, suggest that high-risk drinkers and alcoholic patients selectively pay attention to, interpret, and recall alcohol-related words, and they feel appetitive valence and high arousal when reading them. Our results also indicate a significant negative correlation between AUDIT scores and dominance ratings, indicating that people with greater alcohol consumption feel less in control when reading words that depict alcohol content. This effect, together with previous results, favors approach behavior.

Such findings may have implications for alcohol-related advertisements and therapeutic techniques (Field, Marhe, & Franken, 2014; Gantiva & Flórez-Alarcón, 2015). Alcohol-related advertising and cues appear to have a greater impact on high-risk drinkers and alcoholic patients compared with moderate drinkers (Loeber et al., 2007). Alcohol-related words in advertising may have a greater influence on maintaining and increasing alcohol consumption than on the onset of intake.

The present study has two limitations. First, neither non-drinkers nor alcoholic patients participated in the study. Second, the present study did not assess psychophysiological measures. Future studies should incorporate psychophysiological measures of emotional and motivational responses, with the purpose of increasing the objectivity of the data.

Role of funding sources

Funding for this study was provided by the University of San Buenaventura (Ref. PSI 008-002). University of San Buenaventura, had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

Contributors

The first author designed the study, conducted the statistical analysis and wrote the protocol. The second and third authors conducted literature searches and provided summaries of previous research studies. All authors contributed to and have approved the final manuscript.

Conflict of interest

All authors declare that they have no conflicts of interest.

References

- Aquino, J.M., & Arnell, K.M. (2007). Attention and the processing of words: Dissociating effects of arousal. *Psychonomic Bulletin & Review*, 14, 430–435.
- Arnell, K.M., Killman, K.V., & Fijavz, D. (2007). Blinded by emotion: Target misses follow attentional capture by arousing distractors in RSVP. *Emotion*, 7(3), 465–477. <http://dx.doi.org/10.1037/1528-3542.7.3.465>.
- Bradley, M.M., & Lang, P.J. (1994). Measuring emotion: The self-assessment manikin and the semantic differential. *Journal of Behavior Therapy and Experimental Psychiatry*, 25, 49–59.
- Bradley, M.M., & Lang, P.J. (1999). *Affective Norms for English Words (ANEW): Instruction manual and affective ratings*. Gainesville, FL: Center for Research in Psychophysiology, University of Florida.
- Field, M., Marhe, R., & Franken, I.H. (2014). The clinical relevance of attentional bias in substance use disorders. *CNS Spectrums*, 19(3), 225–230. <http://dx.doi.org/10.1017/S1092852913000321>.
- Fridrici, C., Driessen, M., Wingenfeld, K., Kremer, G., Kissler, J., & Beblo, T. (2014). Investigating biases of attention and memory for alcohol-related and negative words in alcohol-dependents with and without major depression after day-clinic treatment. *Psychiatry Research*, 218, 311–318. <http://dx.doi.org/10.1016/j.psychres.2014.03.04>.
- Gantiva, C., & Flórez-Alarcón, L. (2015). Entrevista motivacional en consumidores excesivos de alcohol: intervenciones efectivas para el aumento en la disposición al cambio. *Psicología: Avances de la Disciplina*, 9(1), 83–91.
- Gendron, M., Lindquist, K.A., Barsalou, L., & Barrett, L.F. (2012). Emotion words shape emotion percepts. *Emotion*, 12(2), 314–325. <http://dx.doi.org/10.1037/a0026007>.
- Harris, C.R., & Pashler, H. (2004). Attention and the processing of emotional words and names: Not so special after all. *Psychological Science*, 15, 171–178.
- Herbert, C., Deusch, R., Sütterlin, S., Kübler, A., & Pauli, P. (2011). Negation as a means for emotion regulation? Startle reflex modulation during processing of negated emotional words. *Cognitive, Affective, & Behavioral Neuroscience*, 11(2), 199–206. <http://dx.doi.org/10.3758/s13415-011-0026-1>.
- Herbert, C., Kissler, J., Junghöfer, M., Peyk, P., & Rockstroh, B. (2006). Processing of emotional adjectives: Evidence from startle EMG and ERPs. *Psychophysiology*, 43(2), 197–206. <http://dx.doi.org/10.1111/j.1469-8986.2006.00385.x>.
- Johnsen, B.H., Laberg, J.C., Cox, W.M., Vaksdal, A., & Hugdahl, K. (1994). Alcoholic subjects' attentional bias in the processing of alcohol-related words. *Psychology of Addictive Behaviors*, 8(2), 111–115.
- Kristensen, C.H., Gomes, C., Justo, A., & Vieira, K. (2011). Brazilian norms for the Affective Norms for English Words. *Trends Psychiatry and Psychotherapy*, 33(3), 135–146.
- Lang, P.J. (2010). Emotion and motivation: Toward consensus definitions and a common research purpose. *Emotion Review*, 2(3), 229–233. <http://dx.doi.org/10.1177/1754073910361984>.
- Loeber, S., Croissant, B., Nakovics, H., Zimmer, A., Georgi, A., Klein, S., et al. (2007). The startle reflex in alcohol-dependent patients: Changes after cognitive-behavioral therapy and predictive validity for drinking behavior. *Psychotherapy and Psychosomatics*, 76(6), 385–390.
- Redondo, J., Fraga, I., Padrón, I., & Comesaña, M. (2007). The Spanish adaptation of ANEW (Affective Norms for English Words). *Behavior Research Methods*, 39, 600–605.
- Soares, A.P., Comesaña, M., Pinheiro, A.P., Simões, A., & Frade, C.S. (2012). The adaptation of the Affective Norms for English words (ANEW) for European Portuguese. *Behavior Research Methods*, 44(1), 256–269. <http://dx.doi.org/10.3758/s13428-011-0131-7>.
- Stormark, K.M., Laberg, J.C., Nordby, H., & Hugdahl, K. (2000). Alcoholics' selective attention to alcohol stimuli: Automated processing? *Journal of Studies on Alcohol*, 61(1), 18–23.
- van Zyl, H., & Meiselman, H.L. (2015). The roles of culture and language in designing emotion lists: Comparing the same language in different English and Spanish speaking countries. *Food Quality and Preference*, 41, 201–213. <http://dx.doi.org/10.1016/j.foodqual.2014.12.00>.
- WHO [World Health Organization] (2001). *AUDIT*. Questionario de Identificación de los Transtornos debidos al Consumo de Alcohol. Pautas para su utilización en Atención Primaria (Available in: http://www.who.int/substance_abuse/activities/en/AUDITmanualSpanish.pdf).
- Williams, J.M.G., Mathews, A., & MacLeod, C. (1996). The emotional Stroop task and psychopathology. *Psychological Bulletin*, 120, 2–24.
- Woud, M.L., Pawelczak, S., Rinck, M., Lindenmeyer, J., Souren, P., Wiers, R.W., et al. (2014). Alcohol-related interpretation bias in alcohol-dependent patients. *Alcoholism: Clinical and Experimental Research*, 38, 1151–1159. <http://dx.doi.org/10.1111/acer.12334>.